SILUAN'S EARLY WARNING SIGNS OF CHILDHOOD CANCER

To improve the survival rate, more children and teens need to be diagnosed at early stages and treated by a Paediatric Oncologist.



"Keeping more than hope alive"





IS FOR SEEK

Medical help early for persistent symptoms.



IS FOR FYF

White spot in the eye, new squint, new blindness, bulging eyeball.



IS FOR LUMP

Abdomen and pelvis, head and neck, limbs, testes and glands.



IS FOR UNEXPLAINED

Prolonged fever over two weeks, loss of weight, fatigue, easy bruising or bleeding.



IS FOR ACHING

Bones, joints, back and easy fractures.



IS FOR NEUROLOGICAL SIGNS

Change in behaviour, balance, walk, milestones. Headaches with early morning vomiting.

HEAD OFFICE

Tel: 086 111 3500 Email: headoffice@choc.org.za

GALITENG SOUTH

Tel: 011 486 1212 Email: receptionjhb@choc.org.za

NORTHERN REGION

Tel: 012 323 3490 Email: pretoria@choc.org.za KWAZULU-NATAL

Ourban Tel: 031 240 2917 Email: dbn@choc.org.z

EASTERN CAPE

East London Tel: 043 748 5315 Email: el@choc.org.za

Port Elizabeth Tel: 041 464 0005 Email: pe@choc.org FREE STATE & KALAHARI

Tel: 051 522 6671 Email: freestate@choc.org.za

WESTERN CAPE & NAMAKWA

Tel: 021 531 0052 Email: wcape@choc.org.za





